arXiv:1604.04934v2 [math.DG] 11 Jul 2016

THE INDEX OF SYMMETRY OF THREE-DIMENSIONAL LIE GROUPS WITH A LEFT-INVARIANT METRIC

SILVIO REGGIANI

Abstract. We determine the index of symmetry of 3-dimensional unimodular Lie groups with a left-invariant metric. In particular, we prove that every 3-dimensional unimodular Lie group admits a left-invariant metric with positive index of symmetry. We also study the geometry of the quotients by the so-called foliation of symmetry, and we explain in what cases the group fibers over a 2-dimensional space of constant curvature.

1. Introduction

The index of symmetry of a Riemannian manifold M is a geometric invariant which measures how far is M from being a symmetric space. Informally, it can be defined as the infimum, over $p \in M$, of the maximum number of independent directions in T_pM arising from Killing fields parallel at p. That is, if the index of symmetry of M is k, then for each $p \in M$ there exist at least k Killing fields $X_1, \dots, X_k$ such that $X_1(p), \dots, X_k(p)$ are linear independent in T_pM and $(\nabla X_i)_p = 0$ for all $i = 1, \dots, k$. So, the index of symmetry of M equals $\dim M$ if and only if M is a symmetric space. This notion appears first in [ORT14] and it turned out to be an interesting way of studying homogeneous spaces. Most of the work was done in the compact case (see [ORT14, BOR15, Pod15]) including the classification of compact spaces with low co-index of symmetry.

The purpose of this article is to compute the index of symmetry of 3-dimensional unimodular Lie groups with a left-invariant metric. Recall that, up to isomorphism, there are six 3-dimensional simply connected, unimodular Lie groups: the abelian Lie group $\mathbb{R}^3$; $\mathrm{SU}(2)$ which is diffeomorphic to a 3-dimensional sphere, the universal covering $\widetilde{\mathrm{SL}}(2,\mathbb{R})$ of the real special linear group of degree 2; the 3-dimensional Heisenberg Lie group H_1; the universal covering $\tilde{E}(2)$ of the group of rigid motions of the euclidean plane; and the group of rigid motions $E(1,1)$ of the Minkowski 2-space. These groups are of a particular importance since most of the Thurston geometries —except $\mathbb{S}^2 \times \mathbb{R}$, $\mathbb{H}^2 \times \mathbb{R}$ and $\mathbb{H}^3$— can be realized as a left-invariant metric on a unimodular 3-dimensional group. Since all the left-invariant metrics on $\mathrm{SU}(2)$ with non-trivial index of symmetry were determined in [BOR15], we direct our attention to the non-compact unimodular Lie groups. As a consequence of our study we can state the following result.

Theorem. *Every 3-dimensional unimodular Lie group admits a left-invariant metric with positive index of symmetry.*

The last part of the article concerns the geometry of the so-called foliation of symmetry. The leaves of the foliation of symmetry $\mathcal{L}$ of the Riemannian manifold

Date: July 12, 2016.

2010 *Mathematics Subject Classification.* 53C30, 53C35.

Key words and phrases. Index of symmetry, unimodular Lie group, distribution of symmetry, naturally reductive space.

Supported by CONICET. Partially supported by ANPCyT and SeCyT-UNR.

M consist of totally geodesic submanifolds which are extrinsically isometric to a symmetric space and of maximal dimension (this is defined precisely in Section 2). We prove that when M is a 3-dimensional unimodular Lie group, the foliation of symmetry induces a Riemannian submersion $M \to M/\mathcal{L}$ if and only if M is $\mathrm{SU}(2)$, $\widetilde{\mathrm{SL}}(2,\mathbb{R})$ or H_1 and the metric is naturally reductive. For the last two cases, we recover the line bundles $\widetilde{\mathrm{SL}}(2,\mathbb{R}) \to \mathbb{H}^2$ over the hyperbolic plane, an the nontrivial vector bundle $H_1 \to \mathbb{R}^2$. It is also worth mentioning that for these naturally reductive metrics, the foliation of symmetry is given by the set of fixed points of the full isotropy group. Thus, these spaces share this important property with compact naturally reductive spaces (see [ORT14]).

Finally, we want to point out that the article includes, in Section 2, a general result which says that the distribution of symmetry of a homogeneous space cannot be of codimension 1. This theorem requires some delicate arguments, including an identification of Killing fields on a Riemannian manifold M with parallel sections of the associated vector bundle $TM \oplus \Lambda^2(TM)$ (see [Kos55, CO08]).

Acknowledgement. The author wants to thank Carlos Olmos for his very useful comments and suggestions on the manuscript, specially the inclusion of Theorem 2.2.

2. Preliminaries

In this section we essentially present the definitions and basic results concerning the index of symmetry of an homogeneous manifold. We refer to [ORT14] and [BOR15] for more details and some structure theory for compact manifolds. Let $M = G/H$ be a Riemannian homogeneous manifold, we say that M has *index of symmetry* $i_{\mathfrak{s}}(M) = k$ if for (any) $p \in M$ there exist Killing fields $X_1, \dots, X_k$ such that $X_1(p), \dots, X_k(p)$ are linear independent in T_pM, $(\nabla X_1)_p = \cdots = (\nabla X_k)_p = 0$ and k is the greatest integer with such properties. In particular, $i_{\mathfrak{s}}(M) = \dim M$ if and only if M is a symmetric space. The so-called *distribution of symmetry* $\mathfrak{s}$ of M is defined as

$$\mathfrak{s}_p = \{v \in T_pM : \text{there exists a Killing field } X \text{ with } X_p = v \text{ and } (\nabla X)_p = 0\}.$$

Note that $\mathfrak{s}$ is G-invariant and moreover, is invariant under the full isometry group $\mathrm{Iso}(M)$ of M. We have that $\mathfrak{s}$ is autoparallel (i.e., integrable with totally geodesic leaves). The integral manifold $L(p)$ of $\mathfrak{s}$ by p is called the *leaf of symmetry* by p. It is known that $L(p)$ is extrinsically isometric to a symmetric space. The *foliation of symmetry* $\mathcal{L}$ of M consists of all the leaves of symmetry of M.

Example 2.1 ([ORT14])**.** If $M = G/H$ is a compact locally irreducible naturally reductive space, and M is not a symmetric space, then the distribution of symmetry of M is the G-invariant distribution induced by the set of fixed vectors of H in the tangent space (via the isotropy representation). In this case, the leaf of symmetry at a given point is a symmetric space of the group type.

In [BOR15] it is proved that $i_{\mathfrak{s}}(M) \leq \dim M - 2$ whenever M is a homogeneous compact space which is not symmetric. We adapt the argument to non-compact spaces and, in particular, to our case of interest (even to non-unimodular 3-dimensional groups).

Theorem 2.2. *Let M be a simply connected homogeneous Riemannian manifold of dimension n, then $i_{\mathfrak{s}}(M) \neq n - 1$.*

The proof of this theorem requires a general framework to deal with Killing fields.

Remark 2.3. Killing fields on a Riemannian manifold M can be regarded as parallel sections of the canonical associated vector bundle $TM \oplus \Lambda^2(TM) \simeq TM \oplus \mathfrak{so}(TM)$, with respect to the connection $\bar{\nabla}$ on $TM \oplus \mathfrak{so}(TM)$ given by

$$\bar{\nabla}_X(Y, B) = (\nabla_X Y - BX, \nabla_X B - R_{X,Y}),$$

where Y is a tangent field on M, B is a skew-symmetric tensor on M of type $(1,1)$ and ∇ is the Levi-Civita connection on M. In fact, if (Y, B) is a parallel section, then $\nabla Y = B$ and so Y is a Killing field on M. Conversely, if Y is a Killing field on M, then its canonical lift $(Y, \nabla Y)$ is a parallel section of $TM \oplus \mathfrak{so}(TM)$, since the derivative of Y satisfies the so-called affine Killing equation $\nabla_X(\nabla Y) = R_{X,Y}$ (see for instance [CO08]).

Lemma 2.4. *Let X, X' be two Killing fields on M with initial conditions at $p \in M$ given by (v, B), $(v', B') \in T_pM \oplus \mathfrak{so}(T_pM)$ respectively. Then $[X, X']$ has initial conditions*

$$(B'v - Bv', R_{v,v'} - [B, B']).$$

Proof. Since $B = (\nabla X)_p$ and $B' = (\nabla X')_p$, the first initial condition holds trivially. For the second one, we use the affine Killing equation (see Remark 2.3) and Bianchi first identity. In fact, if $w \in T_pM$ then

$$\begin{aligned}
\nabla_w[X, X'] &= \nabla_w(\nabla_X X' - \nabla_{X'} X) = \nabla_w(\nabla_X X') - \nabla_w(\nabla_{X'} X) \\
&= (\nabla_w(\nabla X'))_v + \nabla_{\nabla_w X} X' - (\nabla_w(\nabla X))_{v'} - \nabla_{\nabla_w X'} X \\
&= R_{w,v'}v + B'Bw - R_{w,v}v' - BB'w \\
&= R_{v,v'}w - [B, B']w = (R_{v,v'} - [B, B'])w. \qquad \square
\end{aligned}$$

Proof of Theorem 2.2. Assume that the distribution of symmetry of M has dimension $\dim \mathfrak{s} = n - 1$. So, given $p \in M$, we can choose $v \in T_pM$ and a vector field X on M which is always tangent to $\mathfrak{s}$ and such that $\nabla_v X \notin \mathfrak{s}_p$. Recall that $(X, 0)$ is a section of the $\bar{\nabla}$-parallel and flat subbundle $\mathfrak{s} \oplus \mathfrak{so}(\mathfrak{s})$ of the canonical bundle $TM \oplus \mathfrak{so}(TM)$. Since

$$\bar{\nabla}_v(X, 0) = (\nabla_v X, -R_{v,X(p)}) \in \mathfrak{s}_p \oplus \mathfrak{so}(\mathfrak{s}_p),$$

there exists a Killing field Z on M with initial conditions $Z(p) = \nabla_v X$ and $(\nabla Z)_p = -R_{v,X(p)}$. Moreover, we can assume that $Z(p)$ is orthogonal to $\mathfrak{s}_p$. In fact, let w be the orthogonal projection of $\nabla_v X$ to $\mathfrak{s}_p$ and subtract to Z the Killing field with initial conditions $(w, 0)$. We can also assume that v is orthogonal to $\mathfrak{s}_p$, otherwise we subs-tract to Z the Killing field with initial conditions $(0, R_{u,X(p)})$, where u is orthogonal projection of v to $\mathfrak{s}_p$. (Such a Killing field exists, since from Lemma 2.4, the bracket $[U, X']$ of the Killing fields U and X' with initial conditions $(u, 0)$ and $(X(p), 0)$ respectively, has this property.)

We have that Z is always perpendicular to the leaf of symmetry $L(p)$ by p. In fact, let $\bar{Z}$ be the projection to $L(p)$ of the restriction of Z to $L(p)$. Then $\bar{Z}$ is a Killing field of $L(p)$ with initial conditions $(0, (\nabla \bar{Z})_p)$. Since $L(p)$ is totally geodesic, we have that

$$\langle (\nabla \bar{Z})_p v_1, v_2 \rangle = -\langle R_{v,X(p)} v_1, v_2 \rangle = 0$$

for all $v_1, v_2 \in T_pL(p) = \mathfrak{s}_p$. So, the initial conditions of $\bar{Z}$ both vanish and $\bar{Z} = 0$. Therefore Z is always perpendicular to $L(p)$ and moreover, Z must be always perpendicular to the distribution of symmetry $\mathfrak{s}$. If fact, $\mathfrak{s}$ is invariant by isometries and the flow of Z applied to $L(p)$ is open, since $\dim \mathfrak{s} = n - 1$.

We also observe that the Killing field Z, orthogonal to the distributions of symmetry, is unique up to a constant multiple. In fact, let Z' be a Killing field which is always perpendicular to $\mathfrak{s}$ and such that $Z'(p) = 0$. Let Φ'_t be the flow of Z'. Since

$\Phi'_t(p) = p$, we have that $\Phi'_t(L(p)) = L(p)$, since Φ'_t preserves the foliation of symmetry. So, Z' is always tangent to $L(p)$ and thus $Z'|_{L(p)} = 0$. Then $(\nabla Z')_p|_{\mathfrak{s}_p} = 0$ and, since $\mathfrak{s}_p$ has codimension 1, it follows that $(\nabla Z')_p = 0$. This shows that $Z' = 0$, and so Z is unique up to a constant multiple.

It follows from the Killing equation that $\langle \nabla_Z Z, Z\rangle = 0$, and so $\nabla_Z Z$ lies in the distribution of symmetry. Let $\mathfrak{d}$ be the 2-dimensional distribution spanned by Z and $\nabla_Z Z$. We will show that $\mathfrak{d}$ is a parallel distribution. Using this fact, we conclude that M splits as a 2-dimensional symmetric by a $(n-1)$-symmetric space, since $\mathfrak{d} \cap \mathfrak{s} = \mathbb{R}\nabla_Z Z$ is a parallel distribution when restricted to $L(p)$, and this is a contradiction.

In order to show that $\mathfrak{d}$ is a parallel distribution, we note that $\mathfrak{d}$ is invariant under isometries, since Z is unique up to a scalar multiple. This implies that the 1-dimensional distribution $\mathfrak{d} \cap \mathfrak{s}$ is parallel along any leaf of symmetry $L(q)$, $q \in M$. In fact, if U is a Killing field with initial conditions at q, given by $(u, 0)$ with $u \in \mathfrak{s}_q$, then $[U, \nabla_Z Z]$ must be a (functional) multiple of $\nabla_Z Z$, and therefore $\mathfrak{d} \cap \mathfrak{s}$ is parallel along $L(q)$. Since $L(q)$ is totally geodesic and $\mathfrak{d}^\perp \subset \mathfrak{s}$, we conclude that $\mathfrak{d}^\perp$ is autoparallel. Let us prove now that $\mathfrak{d}$ is also autoparallel. Normalize Z and $\nabla_Z Z$ in order to get two unit vector fields Y_1, Y_2 spanning $\mathfrak{d}$. So, Y_1 is orthogonal to $\mathfrak{s}$ and Y_2 lies on $\mathfrak{s}$. Moreover, the integral lines of Y_2 are geodesics (always tangent to the distribution of symmetry), and thus $\nabla_{Y_2} Y_2 = 0$. Since $\nabla_Z Z$ is proportional to Y_2, we also have that $\nabla_{Y_1} Y_1$ lies in $\mathfrak{d}$. Observe now that Y_1 is a unit vector field normal to any totally geodesic hypersurface $L(q)$. So, Y_1 must be parallel along any geodesic of $L(q)$. In particular, $\nabla_{Y_2} Y_1 = 0$. This implies that $\nabla_{Y_1} Y_2 = [Y_1, Y_2]$, which lies in $\mathfrak{d}$. In fact, $[Z, Y_2] = 0$ since $\mathbb{R}Y_2 = \mathfrak{d} \cap \mathfrak{s}$ is invariant under isometries. So $\mathfrak{d}$ and $\mathfrak{d}^\perp$ are both autoparallel and this implies that $\mathfrak{d}$ is a parallel distribution. □

Remark 2.5. We will use repeatedly the well-known Koszul formula in Killing fields (see for instance [BOR15]): if M is a Riemannian manifold and X^*, Y^*, Z^* are Killing fields on M, then

$$\langle \nabla_{X^*} Y^*, Z^*\rangle = \frac{1}{2}\left(\langle [X^*, Y^*], Z^*\rangle + \langle [X^*, Z^*], Y^*\rangle + \langle [Y^*, Z^*], X^*\rangle\right). \tag{2.1}$$

In the case of a Lie group it is also useful the Koszul formula on left-invariant fields: if G is a Lie group endowed with a left-invariant metric and X, Y, Z are left-invariant fields on G, then

$$\langle \nabla_X Y, Z\rangle = \frac{1}{2}\left(\langle [X, Y], Z\rangle - \langle [X, Z], Y\rangle - \langle [Y, Z], X\rangle\right). \tag{2.2}$$

3. The case of the special unitary group SU(2)

Let us consider for the special unitary Lie group SU(2), the basis of its Lie algebra $\mathfrak{su}(2) = \{A \in \mathfrak{gl}(2, \mathbb{C}) : A + A^* = 0\}$ given by

$$X_1 = \frac{1}{2}\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \qquad X_2 = \frac{1}{2}\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \qquad X_3 = \frac{1}{2}\begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix},$$

where $i = \sqrt{-1}$. Any left-invariant metric on SU(2) is identified with a scalar product on $\mathfrak{su}(2)$. Moreover, it follows from [HL09] that any left-invariant metric is automorphically isometric to one of the form $g_{\lambda,\mu,\nu}$, which is represented in the basis X_1, X_2, X_3 by the symmetric positive-definite matrix

$$\begin{pmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}$$

where $\lambda \geq \mu \geq \nu > 0$. In particular, such a metric is bi-invariant if and only if $\lambda = \mu = \nu$. In [BOR15], the left-invariant metrics on $\mathrm{SU}(2)$ with non-trivial index of symmetry are computed. Moreover, this family classifies all the simply connected, compact, homogeneous manifolds with co-index of symmetry equals to 2.

Theorem 3.1 (See [BOR15])**.** *Let g be a left-invariant metric on $\mathrm{SU}(2)$. Then*

1. *$i_{\mathfrak{s}}(\mathrm{SU}(2), g) = 3$ if and only if g is bi-invariant, and in this case $(\mathrm{SU}(2), g)$ is isometric to a round sphere;*
2. *$i_{\mathfrak{s}}(\mathrm{SU}(2), g) = 1$ if and only if, up to isometry and scaling, $g = g_{\lambda,\lambda-1,1}$ with $\lambda > 2$, $g = g_{\lambda,1,1}$ with $\lambda > 1$, or $g = g_{1,1,\nu}$, with $0 < \nu < 1$.*

Remark 3.2. Recall that from [BOR15] (see also [HL12]), all the isometry groups of the metrics of the family $g_{\lambda,\lambda-1,1}$ have dimension 3. So, the symmetric subspace (inside the isometry algebra) for these metrics is spanned for a right-invariant vector field. In contrast, the metrics $g_{\lambda,1,1}$ and $g_{1,1,\nu}$ have all 4-dimensional isometry group, and the symmetric subspaces are never spanned by right-invariant Killing fields except for $g_{2,1,1}$, which correspond to unit tangent bundle of the 2-dimensional sphere of curvature 2 with the Sasaki metric (see [ORT14]).

4. The case of special linear group $\mathrm{SL}(2, \mathbb{R})$

Since the distribution of symmetry lifts to the universal covering, we can perform our calculations on $\mathrm{SL}(2, \mathbb{R})$ instead of $\widetilde{\mathrm{SL}}(2, \mathbb{R})$. Let us consider the basis of the Lie algebra $\mathfrak{sl}(2, \mathbb{R}) = \{A \in \mathfrak{gl}(2, \mathbb{R}) : A + A^t = 0\}$ given by

$$X_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \qquad X_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad X_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

and note that $[X_1, X_2] = 2X_3$, $[X_1, X_3] = -2X_2$ and $[X_2, X_3] = -2X_1$. We have from [HL09] that any left-invariant metric on $\mathrm{SL}(2, \mathbb{R})$ is represented, up to isometric automorphism, in the basis X_1, X_2, X_3 by the symmetric positive-definite matrix

$$\begin{pmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix},$$

where, $\lambda > 0$ and $\mu \geq \nu > 0$. Let us denote by X_1^*, X_2^*, X_3^* the right-invariant vector fields with the same initial conditions as X_1, X_2, X_3. We consider separately two different geometric cases.

4.1. **Case $\mu > \nu$.** Here, by the results in [HL12], the group of isometries of $g_{\lambda,\mu,\nu}$ has dimension 3, and so any Killing field is right-invariant. We are looking for a nontrivial Killing field

$$Y^* = aX_1^* + bX_2^* + cX_3^*$$

such that $(\nabla Y^*)_e = 0$. Taking into account that $[X_i^*, X_j^*] = -[X_i, X_j]^*$ we get that

$$\begin{aligned} [X_1^*, Y^*] &= 2(cX_2^* - bX_3^*), \\ [X_2^*, Y^*] &= 2(cX_1^* + aX_3^*), \\ [X_3^*, Y^*] &= -2(bX_1^* + aX_2^*). \end{aligned} \tag{4.1}$$

We denote, for the sake of simplicity, $g_{\lambda,\mu,\nu} = \langle \cdot, \cdot \rangle$. Then, the condition $(\nabla Y^*)_e = 0$ is equivalent to $\langle \nabla_{X_i^*} Y^*, X_j^* \rangle_e = 0$ for $i < j$. By making use of (2.1) and (4.1), we obtain

$$\begin{aligned} 0 &= 2\langle \nabla_{X_1^*} Y^*, X_2^* \rangle_e \\ &= 2c\langle X_2^*, X_2^* \rangle_e - 2c\langle X_3^*, X_3^* \rangle_e - 2c\langle X_1^*, X_1^* \rangle_e \\ &= 2c(-\lambda + \mu - \nu). \end{aligned}$$

Similarly,

$$\begin{aligned} 0 &= 2\langle \nabla_{X_1^*} Y^*, X_3^* \rangle_e \\ &= -2b\langle X_3^*, X_3^* \rangle_e + 2b\langle X_2^*, X_2^* \rangle_e + 2b\langle X_1^*, X_1^* \rangle_e \\ &= 2b(\lambda + \mu - \nu) \end{aligned}$$

and

$$\begin{aligned} 0 &= 2\langle \nabla_{X_2^*} Y^*, X_3^* \rangle_e \\ &= 2a\langle X_3^*, X_3^* \rangle_e + 2a\langle X_1^*, X_1^* \rangle_e + 2a\langle X_1^*, X_1^* \rangle_e \\ &= 2a(\lambda + \mu + \nu) \end{aligned}$$

So, $(\nabla Y^*)_e = 0$ if and only if

$$2c(-\lambda + \mu - \nu) = 2b(\lambda + \mu - \nu) = 2a(\lambda + \mu + \nu) = 0. \tag{4.2}$$

Since $\mu > 0$, it must be $a = b = 0$ and we have the non-trivial solution $Y^* = cX_3^*$, $c \neq 0$, if and only if $\lambda = \mu - \nu$.

4.2. **Case $\mu = \nu$.** Recall that from [HL12], the dimension of the full isometry group of $g_{\lambda,\mu,\mu}$ is 4. Moreover, the Lie algebra of $\mathrm{Iso}(\mathrm{SL}(2,\mathbb{R}), g_{\lambda,\mu,\mu})$ is $\mathfrak{sl}(2,\mathbb{R}) \rtimes \mathfrak{so}(2)$, where the isotropy algebra $\mathfrak{so}(2)$ acts only on $\mathrm{span}_{\mathbb{R}}\{X_2^*, X_3^*\}$. So, there exists a Killing field X_4^* (the same for all λ, μ) such that $X_4^*(e) = 0$ and

$$[X_1^*, X_4^*] = 0, \qquad [X_2^*, X_4^*] = -X_3^*, \qquad [X_3^*, X_4^*] = X_2^*.$$

Let us denote, as in the previous subsection, $g_{\lambda,\mu,\mu} = \langle \cdot, \cdot \rangle$. Now, we are looking for a Killing field

$$Y^* = aX_1^* + bX_2^* + cX_3^* + dX_4^*$$

such that $(\nabla Y^*)_e = 0$. Recall that with same calculations as in (4.2), we have that $d \neq 0$ if Y^* is nontrivial. Moreover, since $X_4(e) = 0$, we have $(\nabla_{X_i^*} X_4)_e = [X_i^*, X_4^*]_e$. If we call $\tilde{Y}^* = Y^* - dX_4^*$, then $\tilde{Y}^*$ is right-invariant, and the formulas for $\langle \nabla_{X_i^*} \tilde{Y}^*, X_j^* \rangle_e$, $1 \leq i < j \leq 3$, are the same as in the previous subsection taking $\mu = \nu$. So, the equations $\langle \nabla_{X_i^*} Y^*, X_j^* \rangle_e = 0$ become

$$\begin{aligned} 0 &= 2\langle \nabla_{X_1^*} Y^*, X_2^* \rangle_e = -2c\lambda \\ 0 &= 2\langle \nabla_{X_1^*} Y^*, X_3^* \rangle_e = 2b\lambda \\ 0 &= 2\langle \nabla_{X_2^*} Y^*, X_3^* \rangle_e = 2a(\lambda + 2\mu) - 2d\mu. \end{aligned}$$

Thus, $b = c = 0$ and we have a nontrivial solution with $d = a(\lambda + 2\mu)/\mu$. In particular, if we take $a = 1$, then $Y^* = X_1^* + (\frac{\lambda}{\mu} + 2)X_4^*$.

We can summarize the results of this section with the following theorem.

Theorem 4.1. *Let g be a left-invariant metric on $\mathrm{SL}(2,\mathbb{R})$ such that $i_{\mathfrak{s}}(\mathrm{SL}(2,\mathbb{R}), g)$ is nontrivial. Then, up to isometry and scaling, $g = g_{\lambda,\lambda+1,1}$ or $g = g_{\lambda,1,1}$, where $\lambda > 0$.*

5. The case of the Heisenberg Lie group H_1

Let us consider the 3-dimensional Heisenberg Lie group modelled as the differentiable manifold $H_1 = \mathbb{R}^3$ with the group structure

$$(x, y, z)(x', y', z') = (x + x', y + y', z + z' + \tfrac{1}{2}(xy' - yx')).$$

Let us denote the identity element of H_1 by $e = (0, 0, 0)$. Recall that a basis of left-invariant vector field is given by

$$X_1 = \frac{\partial}{\partial x} - \frac{y}{2}\frac{\partial}{\partial z}, \qquad X_2 = \frac{\partial}{\partial y} + \frac{x}{2}\frac{\partial}{\partial z}, \qquad X_3 = \frac{\partial}{\partial z} \tag{5.1}$$

in canonical coordinates (x, y, z). Notice that X_3 spans the center of the Lie algebra $\mathfrak{h}_1$ of H_1 and $[X_1, X_2] = X_3$. It is well-known that there exists only one left-invariant metric on H_1, up to isometry and scaling (see Remark 5.2). So, in order to compute the index of symmetry of H_1 with respect to a left-invariant metric, we can assume that it is endowed with left-invariant metric $\langle \cdot, \cdot \rangle$ that makes (5.1) an orthonormal basis.

Notice that, since H_1 is not a symmetric space, according to Theorem 2.2, its index of symmetry must be 0 or 1. So we look for the solutions of $(\nabla Y^*)_e = 0$, where Y^* is a Killing vector field on H_1. It is well-known that the full isometry group of H_1 has 1-dimensional isotropy group. Moreover, the Lie algebra of the isotropy group is generated for the Killing field

$$X_4^* = -yX_1 + xX_2 - \frac{x^2 + y^2}{2} X_3,$$

and so any Killing vector field can be written as the sum of a right-invariant vector field and a constant multiple of X_4^*. Let us consider the basis of right-invariant vector fields X_1^*, X_2^*, X_3^* given by

$$X_1^* = X_1 + yX_3, \qquad X_2^* = X_2 - xX_3, \qquad X_3^* = X_3.$$

Then we can write the Killing vector field Y^* as a linear combination

$$Y^* = aX_1^* + bX_2^* + cX_3^* + dX_4^* \tag{5.2}$$

for some $a, b, c, d, \in \mathbb{R}$. In order to study the equation $(\nabla Y^*)_e = 0$ it is convenient to use Koszul formula in left-invariant fields. It is easy to verify from (2.2) that

$$\nabla_{X_1} X_1 = \nabla_{X_2} X_2 = \nabla_{X_3} X_3 = 0,$$

$$\nabla_{X_1} X_2 = \frac{1}{2} X_3 = -\nabla_{X_2} X_1$$

and

$$\nabla_{X_3} X_1 = \nabla_{X_1} X_3 = -\frac{1}{2} X_2, \qquad \nabla_{X_3} X_2 = \nabla_{X_2} X_3 = \frac{1}{2} X_1.$$

Assume that Y^* as in (5.2) satisfies $(\nabla Y^*)_e = 0$. We have in particular that

$$0 = (\nabla_{X_3} Y^*)_e = a(\nabla_{X_3} X_1^*)_e + b(\nabla_{X_3} X_2^*)_e$$

since also at the origin we have $0 = [X_3, X_4^*]_e = (\nabla_{X_3} X_4^*)_e$. By replacing X_1^* and X_2^* in terms of left-invariant vector fields in the above equation we get that $0 = (b/2)X_1(e) - (a/2)X_2(e)$ and therefore $a = b = 0$. So, equation (5.2) has now the form $Y^* = cX_3^* + dX_4^*$. Note that neither c nor d can be equal to zero, and then we can assume that $c = 1$. Finally, since $(\nabla_{X_1} X_4^*)_e = X_2(e)$ and $(\nabla_{X_2} X_4^*)_e = -X_1(e)$, we have that

$$(\nabla_{X_1} Y^*)_e = (\nabla_{X_2} Y^*)_e = (\nabla_{X_3} Y^*)_e = 0$$

if and only if $d = -1/2$.

Theorem 5.1. *Let H_1 be the 3-dimensional Heisenberg Lie group endowed with the left-invariant metric such that the basis X_1, X_2, X_3 of $\mathfrak{h}_1$ as in (5.1) is orthonormal. Then H_1 has index of symmetry $i_s(H_1) = 1$. Moreover, the Killing vector field*

$$Y^* = X_3 - \frac{1}{2} X_4^*$$

is parallel at the origin, where $X_4^* = -yX_1 + xX_2 - \frac{1}{2}(x^2 + y^2)X_3$.

As mentioned before, the statement of the above theorem is true for any left-invariant metric on H_1.

Remark 5.2. It is shown in [Lau03] and also in [KTT11], that the only 3-dimensional simply connected Lie groups admitting only one left-invariant metric, up to isometry and scaling, are the abelian Lie group $\mathbb{R}^3$, the simply connected Lie group acting simply transitively on the hyperbolic space $\mathbb{H}^3$, and the 3-dimensional Heisenberg Lie group H_1. For the sake of completeness, we give a proof of this fact for the last case. Keeping the notation of the above paragraphs, let $\langle\cdot,\cdot\rangle'$ be an arbitrary inner product on $\mathfrak{h}_1$. By rescaling the metric we can assume that the left-invariant vector field X_3 has norm one with respect to both of the corresponding left-invariant metrics. Call $\mathfrak{v}$ the vector subspace of $\mathfrak{h}_1$ spanned by X_1, X_2, which is the orthogonal complement to $\mathfrak{z}$ with respect to $\langle\cdot,\cdot\rangle$. Let $\mathfrak{v}'$ be the orthogonal complement to $\mathfrak{z}$ with respect to $\langle\cdot,\cdot\rangle'$ and let X_1', X_2' be an orthonormal basis of $\mathfrak{v}'$. It is easy to see that the linear isometry from $(\mathfrak{h}_1, \langle\cdot,\cdot\rangle)$ onto $(\mathfrak{h}_1, \langle\cdot,\cdot\rangle')$ mapping X_1, X_2, X_3 into X_1', X_2', X_3 respectively, lifts to an isometry from $(H_1, \langle\cdot,\cdot\rangle)$ onto $(H_1, \langle\cdot,\cdot\rangle')$.

Recall that the above argument fails in higher dimensions (see [Vuk15]). In fact, given an inner product $\langle\cdot,\cdot\rangle$ on the $(2n+1)$-dimensional Heisenberg Lie algebra $\mathfrak{h}_n$, we decompose as an orthogonal sum $\mathfrak{h}_n = \mathfrak{v} \oplus \mathfrak{z}$ and for every $Z \in \mathfrak{z}$ we have associated the element $j(Z) \in \mathfrak{so}(\mathfrak{v})$ given by $\langle j(Z)X, Y\rangle = \langle [X,Y], Z\rangle$. Such a map is a geometric invariant. For the standard H-type left-invariant metric on H_n, $j(Z)$ acts on $\mathfrak{v}$ as a multiple of the multiplication by $\sqrt{-1}$, once identified $\mathfrak{v} \simeq \mathbb{C}^n$. But this is not always the case for an arbitrary left-invariant metric when $n > 1$.

6. The case of $\tilde{E}(2)$

We present the universal cover $\tilde{E}(2)$ of the Euclidean group $E(2)$, as the underlying manifold $\mathbb{R}^3$ with the group multiplication given by

$$\begin{aligned}(x,y,z)(x',y',z') &= (x',y',z')\begin{pmatrix}\cos z & \sin z & 0\\ -\sin z & \cos z & 0\\ 0 & 0 & 1\end{pmatrix} + (x,y,z)\\ &= (x + x'\cos z - y'\sin z,\, y + x'\sin z + y'\cos z,\, z + z').\end{aligned}$$

Let us consider the basis of left-invariant fields given by

$$X_1 = \cos z\frac{\partial}{\partial x} + \sin z\frac{\partial}{\partial y}, \qquad X_2 = -\sin z\frac{\partial}{\partial x} + \cos z\frac{\partial}{\partial y}, \qquad X_3 = \frac{\partial}{\partial z}.$$

Such a basis satisfies the bracket relations

$$[X_1, X_2] = 0, \qquad [X_1, X_3] = -X_2, \qquad [X_2, X_3] = X_1$$

and one has the associated right-invariant vector fields

$$X_1^* = \frac{\partial}{\partial x}, \qquad X_2^* = \frac{\partial}{\partial y}, \qquad X_3^* = -y\frac{\partial}{\partial x} + x\frac{\partial}{\partial y} + \frac{\partial}{\partial z}.$$

It is a well-known fact that the left-invariant metric on $\tilde{E}(2)$ that makes of X_1, X_2, X_3 an orthonormal basis is flat (see for instance [Mil76]), and hence isometric to the Euclidean 3-dimensional space. Moreover, one can see that any flat metric on $\tilde{E}(2)$ is obtained from this one rescaling by a positive multiple the metric on the abelian ideal generated by X_1 and X_2. The non-flat left-invariant metrics on $\tilde{E}(2)$ are classified, up to isometric automorphism, in [HL09], and they are represented in the basis X_1, X_2, X_3 by the symmetric positive-defined matrix

$$\begin{pmatrix}1 & 0 & 0\\ 0 & \mu & 0\\ 0 & 0 & \nu\end{pmatrix}$$

where $0 < \mu < 1$ and $\nu > 0$. Denote such a metric by $g_{\mu,\nu}$. In [HL12], the isometry groups of $g_{\mu,\nu}$ are computed and, it particular, it follows that the full isometry

group of $g_{\mu,\nu}$ has always dimension 3. So, in order to compute $i_{\mathfrak{s}}(\tilde{E}(2), g_{\mu,\nu})$, we have to decide if there exists a right-invariant field

$$Y^* = aX_1^* + bX_2^* + cX_3^*$$

such that $(\nabla Y^*)_e = 0$. As usual, we denote $g_{\mu,\nu} = \langle \cdot, \cdot \rangle$. Now, using Koszul formula on Killing fields (2.1) and the brackets

$$[X_1^*, Y^*] = cX_2^*, \qquad [X_2^*, Y^*] = -cX_1^*, \qquad [X_3^*, Y^*] = bX_1^* - aX_2^*$$

we must have

$$\begin{aligned} 0 &= 2\langle \nabla_{X_1^*} Y^*, X_2^* \rangle_e \\ &= c\langle X_2^*, X_2^* \rangle_e + \langle [X_1^*, X_2^*], Y^* \rangle_e + c\langle X_1^*, X_1^* \rangle_e \\ &= c(1+\mu), \end{aligned}$$

and so $c = 0$,

$$\begin{aligned} 0 &= 2\langle \nabla_{X_1^*} Y^*, X_3^* \rangle_e \\ &= c\langle X_2^*, X_3^* \rangle_e + b\langle X_2^*, X_2^* \rangle_e - b\langle X_1^*, X_1^* \rangle_e \\ &= b(\mu - 1), \end{aligned}$$

which implies $b = 0$, and finally

$$\begin{aligned} 0 &= 2\langle \nabla_{X_2^*} Y^*, X_3^* \rangle_e \\ &= -c\langle X_1^*, X_3^* \rangle_e - a\langle X_1^*, X_1^* \rangle_e + a\langle X_2^*, X_2^* \rangle_e \\ &= a(1-\mu) \end{aligned}$$

says that $a = 0$. So, there is not a non-zero Killing field Y^* such that $(\nabla Y^*)_e = 0$ and therefore the index of symmetry of $g_{\mu,\nu}$ is equal to 0 for all $0 < \mu < 1$, $\nu > 0$.

Theorem 6.1. *Let g be a left-invariant metric on $\tilde{E}(2)$. Then g has non-trivial index of symmetry if and only if g is flat.*

7. The case of $E(1,1)$

We present the group $E(1,1)$, of rigid motions of the Minkowski 2-space, as the underlying manifold $\mathbb{R}^3$ endowed with the group multiplication

$$(x, y, z)(x', y', z') = (x + e^z x', y + e^{-z} y', z + z').$$

It is easy to see that the left-invariant fields

$$X_1 = e^z \frac{\partial}{\partial x}, \qquad X_2 = e^{-z} \frac{\partial}{\partial y}, \qquad X_3 = \frac{\partial}{\partial z}$$

form a basis of $\mathfrak{e}(1,1)$ such that

$$[X_1, X_2] = 0, \qquad [X_1, X_3] = -X_1, \qquad [X_2, X_3] = X_2.$$

The right-invariant fields X_i^* such that $X_i^*(e) = X_i(e)$ are given by

$$X_1^* = \frac{\partial}{\partial x}, \qquad X_2^* = \frac{\partial}{\partial y}, \qquad X_3^* = x\frac{\partial}{\partial x} - y\frac{\partial}{\partial y} + \frac{\partial}{\partial z}$$

From the results in [HL12], we have that there exist, up to isometric automorphism, two families of left-invariant metrics on $E(1,1)$, which are represented in the basis X_1, X_2, X_3 by the symmetric positive-definite matrices

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \nu \end{pmatrix}, \qquad \begin{pmatrix} 1 & 1 & 0 \\ 1 & \mu & 0 \\ 0 & 0 & \nu \end{pmatrix}, \tag{7.1}$$

where $\nu > 0$ (in both cases) and $\mu > 1$. We denote these metrics by g_ν and $g_{\mu,\nu}$ respectively. It also follows from [HL12] that any left-invariant metric on $E(1,1)$

has 3-dimensional full isometry group. So, for any left-invariant metric on $E(1,1)$, a generic Killing field has the form

$$Y^* = aX_1^* + bX_2^* + cX_3^*.$$

In order to study the equation $(\nabla Y^*)_e = 0$ it is useful to note that

$$[X_1^*, Y^*] = cX_1^*, \qquad [X_2^*, Y^*] = -cX_2^*, \qquad [X_3^*, Y^*] = -aX_1^* + bX_2^*. \tag{7.2}$$

We distinguish two separate cases according to the metrics above.

7.1. **Case g_ν.** Assume that $(\nabla Y^*)_e = 0$ and denote $g_\nu = \langle \cdot, \cdot \rangle$. From (2.1) and (7.2) we must have

$$\begin{aligned} 0 &= 2\langle \nabla_{X_1^*} Y^*, X_2^* \rangle_e \\ &= c\langle X_1^*, X_2^* \rangle_e + \langle [X_1^*, X_2^*], Y^* \rangle_e + c\langle X_2^*, X_1^* \rangle_e \\ &= 0, \end{aligned}$$

so, no condition is imposed on $c \in \mathbb{R}$. On the other hand,

$$\begin{aligned} 0 &= 2\langle \nabla_{X_1^*} Y^*, X_3^* \rangle_e \\ &= c\langle X_1^*, X_3^* \rangle_e + a\langle X_1^*, X_1^* \rangle_e + a\langle X_1^*, X_1^* \rangle_e \\ &= 2a, \end{aligned}$$

which means $a = 0$, and

$$\begin{aligned} 0 &= 2\langle \nabla_{X_2^*} Y^*, X_3^* \rangle_e \\ &= -c\langle X_2^*, X_3^* \rangle_e - b\langle X_2^*, X_2^* \rangle_e - b\langle X_2^*, X_2^* \rangle_e \\ &= -2b, \end{aligned}$$

so $b = 0$. Thus, we conclude that $Y^* = X_3^*$ is a Killing field parallel at e, for any value of $\nu > 0$.

7.2. **Case $g_{\mu,\nu}$.** Assume again that $(\nabla Y^*)_e = 0$ and denote $g_{\mu,\nu} = \langle \cdot, \cdot \rangle$. This case is slightly different from the previous one since X_1, X_2, X_3 is not an orthogonal basis. More precisely, then non-zero inner products are $\langle X_1, X_1 \rangle = \langle X_1, X_2 \rangle = 1$, $\langle X_2, X_2 \rangle = \mu$ and $\langle X_3, X_3 \rangle = \nu$. Once again, using the Koszul formula for Killing fields and (7.2) we have the equations

$$\begin{aligned} 0 &= 2\langle \nabla_{X_1^*} Y^*, X_2^* \rangle_e \\ &= c\langle X_1^*, X_2^* \rangle_e + \langle [X_1^*, X_2^*], Y^* \rangle_e + c\langle X_2^*, X_1^* \rangle_e \\ &= 2c, \end{aligned}$$

which implies $c = 0$,

$$\begin{aligned} 0 &= 2\langle \nabla_{X_1^*} Y^*, X_3^* \rangle_e \\ &= c\langle X_1^*, X_3^* \rangle_e + \langle X_1^*, Y^* \rangle_e + \langle aX_1^* - bX_2^*, X_1^* \rangle_e \\ &= a + b + a - b = 2a, \end{aligned}$$

so $a = 0$, and similarly

$$\begin{aligned} 0 &= 2\langle \nabla_{X_2^*} Y^*, X_3^* \rangle_e \\ &= -c\langle X_2^*, X_3^* \rangle_e - \langle X_2^*, Y^* \rangle_e + \langle aX_1^* - bX_2^*, X_2^* \rangle_e \\ &= -a - b + a - b = -2b, \end{aligned}$$

shows that $b = 0$. Therefore, all the metrics $g_{\mu,\nu}$ have index of symmetry equal to zero.

We summarize the results of this section as follows.

Theorem 7.1. *Let g be a left-invariant metric on $E(1,1)$. Then g has non-trivial index of symmetry if and only if $g = g_\nu$ (as in (7.1)) for some $\nu > 0$. Moreover, $i_{\mathfrak{s}}(E(1,1), g_\nu) = 1$ and the right-invariant Killing field which at the identity is orthogonal to the abelian ideal generated by X_1, X_2 spans the symmetric subspace.*

8. Quotient by the leaves of symmetry

Let G be a 3-dimensional unimodular Lie group endowed with a left-invariant metric such that $i_{\mathfrak{s}}(G) = 1$, and let us denote the foliation of symmetry by $\mathcal{L}$. Assume that the quotient $G/\mathcal{L}$ admits an $\mathrm{Iso}_0(G)$-invariant metric. If $\dim \mathrm{Iso}_0(G) > 3$, then there exists a Killing field $X \neq 0$ on G acting trivially on $G/\mathcal{L}$, since $\dim G/\mathcal{L} = 2$ and so the action of $\mathrm{Iso}_0(G)$ on $G/\mathcal{L}$ cannot be effective. Such a Killing field is always tangent to $\mathcal{L}$, or equivalently, X generates the distribution of symmetry, $\mathfrak{s} = \mathbb{R}X$. Let us denote by $L(p)$ the leaf of symmetry by $p \in G$, and Φ_t the flow of X. If $u, v \in T_eG$ are orthogonal to the leaf of symmetry at p, that is $u, v \in X(p)^\perp$. Then $d\Phi_t(u)$ and $d\Phi_t(v)$ are orthogonal to $L(p)$ at $\Phi_t(p)$ and moreover, $\langle u, v\rangle = \langle d\Phi_t(u), d\Phi_t(v)\rangle$ for all t. This means that the projection $\pi : G \to G/\mathcal{L}$ is a Riemannian submersion.

Proposition 8.1. *If M is one of the Riemannian manifolds $(\mathrm{SU}(2), g_{\lambda,1,1})$ with $\lambda > 1$, $(\mathrm{SU}(2), g_{1,1,\nu})$ with $0 < \nu < 1$, $(\mathrm{SL}(2,\mathbb{R}), g_{\lambda,1,1})$ with $\lambda > 0$, or H_1 with a left-invariant metric, then $M \to M/\mathcal{L}$ is a Riemannian submersion where $\mathcal{L}$ is the foliation of symmetry and $M/\mathcal{L}$ has an $\mathrm{Iso}_0(M)$-invariant metric.*

Remark 8.2.

(1) It is proved in [BOR15] that the families of metrics $g_{\lambda,1,1}$ ($\lambda > 1$) and $g_{1,1,\nu}$ on $\mathrm{SU}(2)$ correspond to Berger spheres. In this case we have that the Riemannian submersion $\mathrm{SU}(2) \to \mathrm{SU}(2)/\mathcal{L} \simeq \mathbb{S}^2$ fibers the round sphere $\mathbb{S}^2$.

(2) Recall that the metrics $g_{\lambda,1,1}$ ($\lambda > 0$) on $\mathrm{SL}(2,\mathbb{R})$ are naturally reductive (with respect to the full isometry group) and in this case we have that the distribution of symmetry coincides with the $\mathrm{Iso}_0(\mathrm{SL}(2,\mathbb{R}))$-invariant distribution induced by the fixed points of the isotropy. So the foliation of symmetry of $g_{\lambda,1,1}$ consists of closed geodesics and the Riemannian submersion $\mathrm{SL}(2,\mathbb{R}) \to \mathrm{SL}(2,\mathbb{R})/\mathcal{L} \simeq \mathbb{H}^2$ is a circle bundle over the hyperbolic plane.

(3) It is easy to see that the Riemannian submersion $H_1 \to H_1/\mathcal{L}$ corresponds to the nontrivial vector bundle $H_1 \to \mathbb{R}^2$. In fact, one has that the Killing field X_3, in the center of $\mathfrak{h}_1$, generates the distribution of symmetry, and so X_1, X_2 span the horizontal distribution. Then by using O'Neill formula we have that the sectional curvature of $H_1/\mathcal{L}$ vanishes. Recall that, also in this case, the metric on H_1 is naturally reductive and the distribution of symmetry is the $\mathrm{Iso}(H_1)$-invariant distribution induced by the fixed points of the isotropy.

Proof of Proposition 8.1. It follows from the discussion at the beginning of the section and the above remark. □

Finally, assume that $i_{\mathfrak{s}}(G) = 1$ and that the isometry group of G is 3-dimensional. Let $Y \neq 0$ be a left-invariant field such that the associated Killing field Y^* satisfies $(\nabla Y^*)_e = 0$. Then $\mathfrak{s}_e = \mathbb{R}Y_e$ and, since $\mathfrak{s}$ is left-invariant, we have $\mathfrak{s} = \mathbb{R}Y$. Assume that, in addition, the quotient by the foliation of symmetry admits a G-invariant metric such that $G \to G/\mathcal{L}$ is a Riemannian submersion. If X^* is a Killing field such that $X^*|_e$ is orthogonal to Y_e, then X^* must have constant length along the integral curve of Y by the identity. By using this observation (see the next remark) we can show that $(\mathrm{SL}(2,\mathbb{R}), g_{\lambda,\lambda+1,1})$ and $(E(1,1), g_\nu)$ never induce a Riemannian submersion onto their quotients by the leaves of symmetry.

Remark 8.3. (1) Let us consider the case $(\mathrm{SL}(2,\mathbb{R}), g_{\lambda,\lambda+1,1})$. Keeping the notation of the above paragraph and Section 4, we can take $Y^* = X_3^*$ and $X^* = X_1^*$. So,

$$\begin{aligned} |X_1^*(\mathrm{Exp}(tX_3)| &= |\,\mathrm{Exp}(-tX_3)X_1\,\mathrm{Exp}(tX_3)| \\ &= |\cosh(2t)X_1 - \sinh(2t)X_2| \\ &= \lambda\cosh(2t) + (\lambda+1)|\sinh(2t)| \end{aligned}$$

is unbounded.

(2) Analogously for $(E(1,1), g_\nu)$, we can choose $Y^* = X_3^*$ and $X^* = X_1^*$. In this case $|X_1^*(\mathrm{Exp}(tX_3)| = e^{-t}$ which is also non-constant.

References

[BOR15] J. Berndt, C. Olmos, and S. Reggiani, *Compact homogeneous Riemannian manifolds with low co-index of symmetry*, to appear in J. Eur. Math. Soc. (JEMS) (2015).

[CO08] S. Console and C. Olmos, *Curvature invariants, Killing vector fields, connections and cohomogeneity*, Proc. Amer. Math. Soc. **137** (2008), no. 3, 1069–1072.

[HL09] K. Y. Ha and J. B. Lee, *Left invariant metrics and curvatures on simply connected three-dimensional Lie groups*, Math. Nachr. **282** (2009), no. 1, 868–898.

[HL12] ———, *The isometry groups of simply connected* 3*-dimensional unimodular Lie groups*, J. Geom. Phys. **62** (2012), 189–203.

[Kos55] B. Kostant, *Holonomy and the Lie algebra of infinitesimal motions of a Riemannian manifold*, Trans. Amer. Math. Soc. **80** (1955), 528–542.

[KTT11] H. Kodama, A. Takahara, and H. Tamaru, *The space of left-invariant metrics on a Lie group up to isometry and scaling*, Manuscripta Math. **135** (2011), no. 1-2, 229–243.

[Lau03] J. Lauret, *Degenerations of Lie algebras and geometry of Lie groups*, Differential Geom. Appl. **18** (2003), no. 2, 177–194.

[Mil76] J. Milnor, *Curvatures of left invariant metrics on Lie groups*, Adv. Math. **21** (1976), 293–329.

[ORT14] C. Olmos, S. Reggiani, and H. Tamaru, *The index of symmetry of compact naturally reductive spaces*, Math. Z. **277** (2014), no. 3–4, 611–628.

[Pod15] F. Podestá, *The index of symmetry of a flag manifold*, Rev. Mat. Iberoam. **31** (2015), no. 4, 1415–1422.

[Vuk15] S. Vukmirović, *Classification of left-invariant metrics on the Heisenberg group*, J. Geom. Phys. **94** (2015), 72–80.

CONICET and Universidad Nacional de Rosario, Dpto. de Matemática, ECEN-FCEIA, Av. Pellegrini 250, 2000 Rosario, Argentina

E-mail address: reggiani@fceia.unr.edu.ar

URL: http://www.fceia.unr.edu.ar/~reggiani